

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)**ScienceDirect**

Procedia Food Science 5 (2015) 176 – 179

**Procedia**  
Food Science

International 58th Meat Industry Conference “Meat Safety and Quality: Where it goes?”

## Isoflavones - from biotechnology to functional foods

Radmila Markovic<sup>a,\*</sup>, Milan Z. Baltic<sup>a</sup>, Marija Pavlovic<sup>a</sup>, Milica Glisic<sup>a</sup>, Stamen Radulovic<sup>a</sup>, Vesna Djordjevic<sup>b</sup>, Dragan Sefer<sup>a</sup>

<sup>a</sup>*Faculty of Veterinary Medicine, Bulevar oslobođenja 18, 11000 Belgrade, Serbia*<sup>b</sup>*Institute of Meat Hygiene and Technology, Kacanskog 13, 11000 Belgrade, Serbia*

---

### Abstract

Isoflavones are isolated from about 300 plant species and belong to the group of phytoestrogens. Having structural similarity with estrogens, they are related to the estrogen receptors, and may exhibit estrogenic and anti-estrogenic effects. They are present in many plant nutrients (including soy, alfalfa, flax seed). The main sources of isoflavones are legumes and there are known commercial preparations of isoflavones, which are natural, safe and alternative materials that can provide an effect similar to estrogen. These compounds have effects on the health status, production characteristics of the animals (poultry, pigs), and the characteristics of the final product ingredients.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of scientific committee of The 58th International Meat Industry Conference (MeatCon2015)

**Keywords:** isoflavones; feed; foods of animal origin

---

### 1. Introduction

During the past two decades, soy isoflavones (commonly genistein, daidzein and equol) have gained significant attention from the scientific community, primarily due to their beneficial effects in the treatment of ageing symptoms in both genders<sup>1</sup>. Namely, some positive outcome was noticed in the soy isoflavone treatment of breast, ovary and prostate cancer, and then osteoporosis, cardiovascular disease, diabetes, asthma, Kawasaki syndrome,

---

\* Corresponding author. Tel.: +38164-409-98-25; fax: +38111-2685-951.

E-mail address: [radmilam@vet.bg.ac.rs](mailto:radmilam@vet.bg.ac.rs)

menopausal symptoms, as well as psychological symptoms associated with aging<sup>2,3,4,5</sup>. Some isoflavones (genistein, daidzein), are considered to be natural selective estrogen receptor modulator for molecules binding to estrogen receptors and have selective activity towards different types of tissue for use in the prevention of treatment of certain diseases<sup>6</sup>. The amounts consumed isoflavones (ISF) foods differ. In Japan, where traditional food is consumed, the daily intake of isoflavones is 100 mg genistein and daidzein. In contrast, in America only about 1-3 mg of isoflavones is the average daily intake<sup>7</sup>.

All of these effects depend on many factors, such as duration intake of isoflavones, their form (conjugated or not) and the degree of absorption, the dose, the individual characteristics - especially the concentration of endogenous estrogen. All this must be known when examining the effects of isoflavones on the health status of people, and therefore, their effects remain somewhat unclear. There are numerous positive effects in many hormone-dependent conditions, but there are many epidemiological studies that indicate harmful effects, especially in children and the development of some types of cancer.

## 2. Sources, use and effects of isoflavones

The best sources of natural isoflavones are legumes, particularly soy. In much smaller quantities, isoflavones can be found in foods of animal origin (meat, milk, eggs, fish)<sup>8,9</sup>. Isoflavones (daidzein, genistein and glycitein) are flavonoids made of two phenolic rings bonded to a carbon triple bond. Special attention was drawn when Kuiper *et al.*<sup>10,11</sup> found that genistein shows affinity for estrogen receptors, whereas the daidzein activity was significantly less (forty times). Isoflavones in nutrients may be present in four different chemical forms: non-conjugated, conjugated glucose (glycosidic form), acetyl glucoside and malonyl glucoside<sup>12</sup>.

Recently, commercial preparations of isoflavones have become very popular; they are used as food for humans and animals (broilers, hens, pigs) in order to obtain safe natural alternative compounds that will take over the role of estrogen. Results of some studies suggest that isoflavones have great opportunities for used in food for people and animals, primarily preventive and therapeutic for many diseases.

Soy isoflavones are of interest and research prominence due to the abundance of these foods in the human diet, but on the other hand, testing the potential transmission of these compounds and their active metabolites from feed to the tissues and organs of animals is of great importance, ie. in foods of animal origin. This could enable increased intake of isoflavones, and result in production of functional foods, such is the case with many other supplements (e.g. Selenium, Conjugated Linoleic Acid)<sup>13,14,15</sup>.

In experiments on broilers, addition of up to 80 mg of isoflavones per kg of feed over three weeks (from day 43 to 63), had a significant impact on the growth of body weight (10 or 20 mg of ISF/kg increased the growth by 13.6 and 16.2%, respectively, and increased the consumption of feed by 7.37% and 11.2%, respectively. Addition of 10 mg of ISF/kg improved the feed conversion by 5.5% during the fattening (42-63 days)<sup>16,17</sup>.

The effects of isoflavones from feed to pig production performances show variability. Cook<sup>18</sup> reported that supplemental ISF (containing the isoflavones genistein, daidzein, and glycitein) increased ADG (average daily gain) in gilts from 6 to 30 kg of BW (body weight) and increased the percentage of mostly red muscle mass, but did not affect the white muscle and daily intake. However, in a second study, Cook<sup>18</sup> reported that dietary genistein did not affect ADG in castrated animals from 5 to 28 kg of BW. It was found that soy isoflavones led to an increase in length, weight and leanness body in male fattening pigs<sup>19</sup>. It appears that the reduced percentage of fat, and growing share of muscle tissue in the meat when isoflavones are added to feed, are more pronounced in males than females. The explanation lies in the hormonal activity of isoflavones, to which males are more responsive than females.

Soy isoflavones affect the meat quality of broilers. Supplementation with ISF 40 mg/kg in the diet increased the water binding capacity to 17.24%, and amounts of 20 or 40 mg/kg significantly increased the pH of the meat. Adding 40 or 80 mg of ISF/kg increased a\* value (redness) and reduced L\* value (lightness) of meat<sup>16</sup>.

Isoflavones in the feed of laying hens significantly affected egg production, egg weight and feed conversion. Shi *et al.*<sup>20</sup> have shown that isoflavones added as supplements in the diet of laying hens during the later period of laying, after the peak production, significantly affect egg production, egg weight and feed conversion. It has been shown that these compounds are involved in calcium metabolism, thus leading to a decrease in intracellular calcium concentration in osteoclasts, which leads to an increase in the amount of calcium for shell formation. Therefore, the strength of the shell, shell thickness, shell percentage ratio of the mass and the mass of the egg exhibit linear-

dependent response to increasing concentrations of isoflavones in feed.

Notable is the effect of these compounds, especially daidzein, on the antioxidant status of poultry and eggs; it is considered that, as a consequence of isoflavones, the activity of antioxidant enzymes increases, and as a result an increase is seen in the content of superoxide dismutase (SOD), catalase and glutathione peroxidase (GSH-Px) in blood plasma, breast meat and eggs<sup>21</sup>. The content of lipid peroxidase in the egg yolk, liver and plasma significantly decreased if the feed contained added daidzein (40 mg/kg), while SOD and GSH-Px remained relatively stable<sup>22</sup>. Effects on the production performance may depend on the type of isoflavone, so it is considered that glycetein has three times higher effects compared to genistein. Because of this, the effects of isoflavones on production traits can be variable.

### 3. Conclusion

These compounds may exhibit many positive effects on human health, and animal production properties, but also on the qualitative properties of foods. So, they have a positive effect on growth and leanness in broilers and pigs, while in laying hens, they increased the number and weight of eggs. Also, isoflavones increased the activity of antioxidant enzymes, and improved the antioxidant status of muscle cells and egg yolks. They are deposited in the yolk, and eggs can be a source of these compounds for people, giving rise to the possibilities for production of so-called, designer egg, ie. eggs enriched with isoflavones as specific active ingredients. Monitoring the presence of these substances in foods leads to increased safety of consumers, but also increases consumer awareness of their content in foods, as active substance exhibiting a certain degree of physiological effects in the human body, as well as in animals. It is certain that a definitive ratification of the conditions and amount of use of isoflavones in food for humans and animals will be produced in the future, preceded by numerous investigations.

### Acknowledgement

This paper was supported by the Ministry of Education, Science and Technological Development, Republic of Serbia, Project “Selected biological hazards to the safety / quality of food of animal origin and the control measures from farm to consumer” (31034).

### References

1. Ajdzanovic V, Medigovic I, Pantelic J, Milosevic V. Soy isoflavones and cellular mechanics. *J Bioenerg Biomembr* 2014;**46**: 99-107.
2. Casini ML, Marelli G, Papaleo E, Ferrari A, D'Ambrosio F, Unfer V. Psychological assesment of the effects of treatment with phytoestrogens on postmenopausal women: a randomized, double-blind, crossover, placebo-controlled study. *Fertil Steril* 2006;**85**:972-8.
3. Messina M. Insights gained from 20 years of soy research. *J Nutr* 2010;**140**:2289-95.
4. Andres S, Abraham K, Appel KE, Lampen A. Risks and benefits of dietary isoflavones for cancer. *Crit Rev Toxicol* 2011;**41**:463-506.
5. Wang Qinglu, Ge Xiaoyue, Tian Xuewen, Zhang Yujun, Zhang Jie, Zhang Pingping. Soy osoflavone: The multipurpose phytochemical (Review). *Biomedical Reports* 2013:697-701.
6. Oseni T, Patel R, Pyle J, Jordan VC. Selective estrogen receptor modulators and phytoestrogens. *Planta Medica* 2008;**74**:1656-65.
7. Barnes S. The Biochemistry, Chemistry and Physiology of the Isoflavones in Soybeans and their Food Products. *Lymphat Res Biol* 2010;**8**:89-98.
8. Kariyil BJ. Phytoestrogens in Animal Origin Foods. *Veterinary World* 2010;**3**:43-5.
9. Kuhnle GC, Gunter, Caterina Dell Aquila, Sue MAspinall, Shirley A Runswick, Angela A Mulligan, Sheila A Bingham, Phytoestrogen Content of Foods of Origin: Dairy Products, Eggs, Meat, Fish, and Seafood. *J Agric Food Chem* 2008; **56**:10099-104.
10. Kuiper GGJM, Enmark E, Peltto-Huikko M, Nilsson S, Gustafsson J-Å. Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci USA* 1996;**93**:5925–30.
11. Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Häggblad J, Nilsson S, Gustafsson JA. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors  $\alpha$  and  $\beta$ . *Endocrinology* 1997;**138**:863–70.
12. Gasperevic-Ivanek V. Fitoestrogeni. *Medix* 2003;**9**:50:90-4.
13. Baltic ZM, Markovic R, Djordjevic V. Nutrition and meat quality. *Tehn mesa* 2011;**52**:154-9.
14. Markovic R, Djordjevic V, Baltic ZM. The importance of conjugated linoleic acid for meat quality. Proceedings International 57th meat industry conference. *Meat and meat products- perspectives of sustainable production*. Belgrade: 10-12th june; 2013. p. 17-23.
15. Markovic R, Todorovic M, Pantic S, Baltic ZM, Ivanovic J, Jakic-Dimic D, Sefer D, Petrujkic B, Radulovic S. Effect of the diet on improvement of the fatty acid composition of pig meat. First International Symposium of Veterinary Medicine “One Health-New Challenges“. Vrdnik: Proceedings; 21-23th May; 2015. p. 148-56.

16. Jiang ZY, Jiang SQ, Lin YC, Xi PB, Yu DQ, Wu TX. Effects of Soybean Isoflavone on Growth Performance, Meat Quality, and Antioxidation in Male Broilers. *Poultry Science* 2007;**86**:1356–1362.
17. Cassidy A. Potential risks and benefits of phytoestrogenrich diets. *Int J Vitam Nutr Res* 2003;**73**:120–6.
18. Cook DR. The effect of dietary soybean isoflavones on the rate and efficiency of growth and carcass muscle content in pigs and rats, PhD Diss. Iowa State Univ: Ames; 1998.
19. Payne R, Bidner T, Southern L, Geaghan J. Effects of dietary soy isoflavones on growth, carcass traits, and meat quality in growing-finishing pigs. *J Anim Sci* 2001;**79**:1230–39.
20. Shi SR, Gu H, Chang LL, Wang ZY, Tong HB, Zou JM. Safety evaluation of daidzain in laying hens: Part I. Effects on laying performance, clinical blood parameters, and organs development. *Food Chem Toxicol* 2013;**55**:684-88.
21. Cai J, Gu H, Shi S, Tong H. Effects of High-Dose Daidzein on Laying Performance, Egg Quality and Antioxidation in Laying Hens. *J Poult Sci* 2013;**50**:237-41.
22. Yin JD, Qi GH, Zhang P, Huo QG. Effect of dietary daidzein on egg cholesterol content and its antioxidant property in layers. *Sci Agricultura Sinica* 2004;**37**:756-61.